

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>5</sup> : A61K 39/00, 39/12, G01N 33/569 G01N 33/577, C12N 7/00, 7/02 C12N 7/04	A1	(11) International Publication Number: <b>WO 93/03760</b> (43) International Publication Date: 4 March 1993 (04.03.93)
--	----	---

(21) International Application Number: PCT/US92/06873

(22) International Filing Date: 17 August 1992 (17.08.92)

## (30) Priority data:

749,839	26 August 1991 (26.08.91)	US
760,713	16 September 1991 (16.09.91)	US
860,444	30 March 1992 (30.03.92)	US


(71)(72) Applicants and Inventors: COLLINS, James, Edward [US/US]; 2658 Riveria Drive South, White Bear Lake, MN 55110 (US). BENFIELD, David, Allen [US/US]; 1509-12th Street South, Brookings, SD 57006 (US). CHLADEK, Danny, W. [US/US]; 19 Evergreen Terrace, St. Joseph, MO 64505 (US). HARRIS, Louis, L. [US/US]; 402 Tanglewood Drive, St. Joseph, MO 64505 (US). GORCYCA, David, E. [US/US]; 2408 Elephant Trail, St. Joseph, MO 64506 (US).

(74) Agent: HAMRE, Curtis, B.; Merchant, Gould, Smith, Edell, Welter & Schmidt, 3100 Norwest Center, 90 South Seventh Street, Minneapolis, MN 55402 (US).

(81) Designated States: AT, AU, BB, BG, BR, CA, CH, CS, DE, DK, ES, FI, GB, HU, JP, KP, KR, LK, LU, MG, MN, MW, NL, NO, PL, RO, RU, SD, SE, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, SN, TD, TG).

**Published***With international search report.*

(54) Title: SIRS VACCINE AND DIAGNOSIS METHOD



(57) Abstract

The invention includes a vaccine and sera for treatment of Mystery Swine Disease (MSD), a method for producing the vaccine, methods for diagnosis of MSD, a viral agent that will mimic "mystery swine disease" and antibodies to the viral agent useful in diagnosis and treatment of MSD. The serum contains mammalian antibodies which are effective in treating MSD.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCI on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FI	Finland	MN	Mongolia
AU	Australia	FR	France	MR	Mauritania
BB	Barbados	GA	Gabon	MW	Malawi
BE	Belgium	GB	United Kingdom	NL	Netherlands
BF	Burkina Faso	GN	Guinea	NO	Norway
BG	Bulgaria	GR	Greece	NZ	New Zealand
BJ	Benin	HU	Hungary	PL	Poland
BR	Brazil	IE	Ireland	PT	Portugal
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	RU	Russian Federation
CG	Congo	KP	Democratic People's Republic of Korea	SD	Sudan
CH	Switzerland	KR	Republic of Korea	SE	Sweden
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovak Republic
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CS	Czechoslovakia	LU	Luxembourg	SU	Soviet Union
CZ	Czech Republic	MC	Monaco	TD	Chad
DE	Germany	MG	Madagascar	TG	Togo
DK	Denmark	MI	Mali	UA	Ukraine
ES	Spain			US	United States of America

-1-

## SIRS VACCINE AND DIAGNOSIS METHOD

5

BACKGROUND OF THE INVENTION

Since 1987, the swine-producing industry has been subjected to a devastating epidemic of an unknown disease, often referred to as "Mystery Swine Disease" [MSD, more recently referred to as "Swine Infertility and Respiratory Syndrome (SIRS)], because researchers have been unable to identify the causative agent. MSD has affected hundreds of thousands of swine throughout North America and Europe. Once one pig is infected with MSD, that one pig can spread the MSD to an entire herd within three to seven days. From 1987 to 1991, the swine industry has lost millions of dollars in revenue as a result of MSD. A recent study estimates that MSD causes a financial loss between \$250 and \$500 per inventoried sow.

MSD causes multiple symptoms in swine. The first symptom of MSD in a breeding herd of swine is usually anorexia and mild pyrexia. In addition, the herd animals may exhibit bluish discolorations in their skin, especially in their ears, teats, snout, and the frontal portions of their necks and shoulders. The affected skin may become irreparably damaged. However, the most devastating symptom of MSD is the reproductive failure that occurs in a breeding herd of swine. MSD causes sows to bear stillborn piglets; undersized, weak piglets with respiratory distress; or piglets which die before they are weaned. Other reproductive symptoms caused by MSD include early farrowing of piglets, a decrease in conception rates, failure in some sows to cycle, and a reduction in the total number of piglets found in a litter. It has been estimated that the number of pigs lost from reproductive failure is about 10 to 15 percent of the annual production of pigs.

Research has been directed toward isolating the causative agent of MSD. A number of potential bacterial

-2-

pathogens have been isolated. However, the types of potential bacterial pathogens have varied between swine-producing farms. Viral investigation has included fluorescent antibody examination, electron microscopic investigation, and serology. These methods have failed to locate the causative agent of MSD. As a result, no one has yet developed a vaccine which can be used to treat MSD in the swine population.

Therefore, it is an objective of the invention to provide a vaccine and sera which, when administered to a breeding swine herd, will reduce the presence of MSD in their population. Another object is to provide a method of treating a population of swine with the vaccine to eradicate MSD from the swine population. Yet another object is to provide a method for diagnosis of MSD.

#### SUMMARY OF THE INVENTION

These and other objects are achieved by the present invention which is directed to a vaccine and sera for prevention and treatment of mystery swine disease and to a method for its diagnosis in swine.

The vaccine is derived from an infectious agent that will infect swine with mystery swine disease (MSD). The infectious agent is obtained from an inoculum of processed tissue of swine infected with the disease, preferably lung tissue. Preferably, the infectious agent is the product of an in vitro mammalian cell culture such as a simian cell line infected with the inoculum of the infected swine tissue. Preferably, the inoculum contains biological particles no greater than about 1.0 micron in size, more preferably 0.5 micron, most preferably no greater than 0.2 micron. It is also preferable that the inoculum has been neutralized with antibodies to common swine diseases.

According to the present invention, a tissue homogenate obtained from piglets in SIRS-affected herds

-3-

consistently reproduced the respiratory and reproductive forms of SIRS when intranasally inoculated in gnotobiotic piglets and pregnant sows. Gnotobiotic piglets so inoculated with either unfiltered or filtered (0.45,  
5 0.22, or 0.1  $\mu$ m) inoculum became anorectic and developed microscopic lung lesions similar to lesions seen in SIRS-affected herds. The same inoculum also caused reproductive effects identical to those seen in SIRS-affected herds. A viral agent has been recovered from  
10 the tissue homogenate. The viral agent causes a disease that mimics SIRS in piglets and pregnant sows. The viral agent has not yet been classified. However, the viral agent is a fastidious, non-hemagglutinating enveloped RNA virus. A viral agent causing SIRS has been deposited on  
15 July 18, 1991 with the American Type Culture Collection in Rockville, Maryland under the accession number ATCC VR-2332.

The serum for treatment of infected swine carries mammalian antibodies to the MSD. It is obtained  
20 from the blood plasma of a mammal (non-swine and swine) pre-treated with the above-described infectious agent.

Alternatively, the serum is formulated from monoclonal antibodies to MSD produced by hybridoma methods.

25 The method for diagnosis of MSD is based upon the use of immunospecific antibodies for MSD. The method calls for combination of a filtered homogenate of a lung biopsy sample or a biopsy sample or similar samples (homogenate or biopsy) from other tissue and the  
30 immunospecific antibodies followed by application of a known detection technique for the conjugate formed by this combination. Immobilization or precipitation of the conjugate and application of such detection techniques as ELISA; RIA; Southern, Northern, Western Blots and the  
35 like will diagnose MSD.

-4-

According to the present invention, therapeutic and diagnostic methods employing antibodies to MSD involve monoclonal antibodies (e.g., IgG or IgM) to the above-described fastidious, non-hemagglutinating enveloped RNA virus. Exemplary antibodies include SDOW 12 and SDOW 17, deposited with the American Type Culture Collection on March 27, 1992 with accession numbers HB 10996 and HB 10997, respectively).

10

#### BRIEF DESCRIPTION

Figure 1 shows the cytopathic effects observed with SIRS virus VR-2332. Figure 1A: Non-infected, unstained cell monolayer. Figure 1B: Infected monolayer with small granular rounded and degenerating cells observed three days post-innoculation with the 6th passage of the SIRS virus.

Figure 2 shows the direct immunofluorescence staining of SIRS virus-infected MA-104 cells.

Figure 2A: Non-infected cell monolayer. Figure 2B: Infected cells with intense, often granular cytoplasmic fluorescence observed three days post-innoculation.

Figure 3 shows the density gradient profile of SIRS virus purified on CsCl density gradients. Peak virus infectivity occurs at 1.18-1.19 g/ml.

Figure 4 shows an electron micrograph of virus particles observed in CsCl gradient fractions of density 1.18-1.19 g/ml. Figure 4A: These four particles are spherical, 60-65 nm in diameter. Two particles are "empty", showing electron-dense core (arrows), and the other two particles are complete. Bar = 100 nm.

Figure 4B: Immuno-gold electron microscopy of SIRS virus with hyperimmune rabbit sera and anti-rabbit IgG labeled with gold particles. Note presence of core particle approximately 25-30 nm in diameter within the virion.

Bar = 50 nm.

-5-

Figure 5 shows the temperature stability of SIRS virus at 4°C (open triangles), 37°C (open circles), and 56°C (closed circles).

5

#### DETAILED DESCRIPTION OF THE INVENTION

Determination of the cause of Mystery Swine Disease (MSD) has been difficult. According to the present invention, however, the isolation and growth of the infectious agent causing MSD has been achieved. As  
10 used herein, "infectious agent" refers to a virus capable of causing swine infertility and respiratory syndrome. More specifically, the infectious agent is a fastidious, non-hemagglutinating enveloped RNA virus and zoopathogenic mutants thereof capable of causing swine  
15 infertility and respiratory disease in swine. The isolation of the infectious agent is a major breakthrough and discovery. It enables the production of vaccines, antibody sera for treatment of infected swine, and diagnostic methods.

20 The vaccine is composed of an inactivated or attenuated MSD infectious agent, derived from an inoculum processed from infected swine lung tissue or other swine tissue exhibiting the characteristic lesions of MSD. Functional derivatives of infectious agent, including  
25 subunit, vector, recombinant, and synthetic peptide vaccines, or the like, are also envisioned. A multi-step procedure is utilized in developing the MSD vaccine. The MSD infectious agent is first obtained as an inoculum by separation and isolation from infected swine tissue,  
30 preferably the lung tissue. The MSD infectious agent is then treated using known vaccinological techniques to form a vaccine against MSD.

The MSD infectious agent is preferably isolated as an inoculate from lung tissue of pigs which exhibit  
35 rapid breathing due to the MSD (other tissue such as fetal tissue may also be used to recover virus). Such

-6-

pigs are destroyed and their lung tissue removed. The lung tissue is then microscopically examined for thickened alveolar septae caused by the presence of macrophages, degenerating cells, and debris in alveolar spaces. These characteristics indicate the presence of the MSD infectious agent. Other swine tissue exhibiting lesions of this sort may also be used to isolate the MSD infectious agent.

The lung or other swine tissue is then homogenized with a pharmaceutically acceptable aqueous solution (such as physiological saline, Ringers solution, Hank's Balanced Salt Solution, Minimum Essential Medium, and the like) such that the tissue comprises 10 percent weight/volume amount of the homogenate. The homogenate is then passed through filters with pore diameters in the 0.05 to 10 micron range, preferably through a series of 0.45, 0.2 and 0.1 micron filters, to produce a filtered homogenate containing the MSD infectious agent. As a result, the filtered homogenate contains biological particles having a size no greater than about 1.0 micron, preferably no greater than about 0.2 to 0.1 micron. The filtered homogenate can then be mixed with Freund's incomplete adjuvant so that the production of antibodies can be stimulated upon injection into a mammal. This mixture can be used as an inoculum for development of MSD in swine or further study of the MSD infectious agent.

After obtaining a filtered homogenate containing the infectious agent, the infectious agent can be inactivated or killed by treatment of the filtered homogenate with a standard chemical inactivating agent such as an aldehyde reagent including formalin, acetaldehyde and the like; reactive acidic alcohols including cresol, phenol and the like; acids such as benzoic acid, benzene sulfonic acid and the like; lactones such as beta propiolactone and caprolactone; and activated lactams, carbodiimides and carbonyl

-7-

diheteroaromatic compounds such as carbonyl diimidazole. Irradiation such as with ultraviolet and gamma irradiation can also be used to inactivate or kill the infectious agent. Alternatively, the infectious agent  
5 can be attenuated by its repeated growth in cell culture from non-swine mammal or avian origin so that the ability of the infectious agent to virulently reproduce is lost. The details of the cell culture attenuation technique are given below.

10           The killed or attenuated infectious agent is then diluted to an appropriate titer by addition of a diluent adjuvant solution for stimulation of immune response. The titration is accomplished by measurement against MSD antibody in an immunologic test such as an  
15 ELISA, RIA, IFA or enzyme substrate detection test as described below.

To produce a purified form of the infectious agent, the filtered homogenate described above can be inoculated into a series of in vitro cell preparations.  
20 Cell preparations with mammalian organ cells such as kidney, liver, heart and brain, lung, spleen, testicle, turbinate, white and red blood cells and lymph node, as well as insect and avian embryo preparations can be used. Culture media suitable for these cell preparations  
25 include those supporting mammalian cell growth such as fetal calf serum and agar, blood infusion agar, brain-heart infusion glucose broth and agar and the like. Preferably the mammalian cells are monkey kidney cells, most preferably African green monkey kidney embryonic  
30 cells -- monkey kidney cell line (MA-104).

After inoculating the cell preparation with the filtered homogenate and growing the culture, individual clumps of cultured cells are harvested and reintroduced into sterile culture medium with cells. The culture  
35 fluid from the final culture of the series provides the purified form of the virulent infectious agent. Also,

-8-

after a series of repeated harvests have been made, the culture can be grown, the culture fluid collected and the fluid used as an inoculum for a culture of a different cellular species. In this fashion, the infective agent  
5 can be attenuated such that the culture fluid from the differing species culture provides the purified form of the attenuated infectious agent.

Polyclonal antibody sera can be produced through use of the infectious agent as an antigenic  
10 substance to raise an immune response in mammals. The culture fluid or inoculum prepared as described above can be administered with a stimulating adjuvant to a non-swine mammal such as a horse, goat, mouse or rabbit. After repeated challenge, portions of blood serum can be  
15 removed and antigenically purified using immobilized antibodies to those disease specific antibodies typically found in the serum of the bled animal. Further treatment of the semi-purified serum by chromatography on, for example, a saccharide gel column with physiological  
20 saline and collection of proteinaceous components of molecular weight at least 10,000 provides a purified polyclonal sera for use in treatment.

Monoclonal antibody sera can be produced by the hybridoma technique. After immunization of a mouse, pig,  
25 rat, rabbit or other appropriate species with MSD containing cell culture lysate or gradient-purified MSD as described above, the spleen of the animal can be removed and converted into a whole cell preparation. Following the method of Kohler and Milstein (Kohler et  
30 al., Nature, 256, 495-97 (1975)), the immune cells from the spleen cell preparation can be fused with myeloma cells to produce hybridomas. Culturation of the hybridomas and testing the culture fluid against the fluid or inoculum carrying the infectious agent allows  
35 isolation of the hybridoma culture producing monoclonal antibodies to the MSD infectious agent. Introduction of

-9-

the hybridoma into the peritoneum of the host species will produce a peritoneal growth of the hybridoma. Collection of the ascites fluid yields body fluid containing the monoclonal antibody to the infectious agent. Also, cell culture supernatant from the hybridoma cell culture can be used. Preferably the monoclonal antibody is produced by a murine derived hybrid cell line wherein the antibody is an IgG or IgM type immunoglobulin. Example monoclonal antibodies to the infectious agent for SIRS are monoclonal antibody SDOW 12 and SDOW 17. In addition to uses discussed elsewhere in this application, monoclonal antibodies according to the present invention can be employed in various diagnostic and therapeutic compositions and methods, including passive immunization and anti-idiotypic vaccine preparation.

The vaccine of the present invention is capable of preventing and curing MSD infections found in the swine population. For effective prophylactic and anti-infectious use in vivo, the MSD vaccine contains killed or attenuated MSD infectious agent and may be administered alone or in combination with a pharmaceutical carrier that is compatible with swine. The vaccine may be delivered orally, parenterally, intranasally or intravenously. Factors bearing on the vaccine dosage include, for example, the age, weight, and level of maternal antibody of the infected pig. The range of a given dose is  $10^3$  to  $10^7$  Tissue Culture Infective Dose 50 per ml, preferably given in 1 ml to 5 ml doses. The vaccine doses should be applied over about 14 to 28 days to ensure that the pig has developed an immunity to the MSD infection.

The MSD vaccine can be administered in a variety of different dosage forms. An aqueous medium containing the killed or attenuated MSD infectious agent may be desiccated and combined with pharmaceutically

-10-

acceptable inert excipients and buffering agents such as lactose, starch, calcium carbonate, sodium citrate formed into tablets, capsules and the like. These combinations may also be formed into a powder or suspended in an aqueous solution such that these powders and/or solutions can be added to animal feed or to the animals' drinking water. These MSD vaccine powders or solutions can be suitably sweetened or flavored by various known agents to promote the uptake of the vaccine orally by the pig.

For purposes of parenteral administration, the killed or attenuated MSD infectious agent can be combined with pharmaceutically acceptable carrier(s) well known in the art such as saline solution, water, propylene glycol, etc. In this form, the vaccine can be parenterally, intranasally, and orally applied by well-known methods known in the art of veterinary medicine. The MSD vaccine can also be administered intravenously by syringe. In this form, the MSD vaccine is combined with pharmaceutically acceptable aqueous carrier(s) such as a saline solution. The parenteral and intravenous formulations of MSD vaccine may also include emulsifying and/or suspending agents as well, together with pharmaceutically acceptable diluent to control the delivery and the dose amount of the MSD vaccine.

The method for diagnosis of MSD is carried out with the polyclonal or monoclonal antibody sera described above. Either the antibody sera or the biopsied tissue homogenate may be immobilized by contact with a polystyrene surface or with a surface of another polymer for immobilizing protein. The other of the antibody sera and homogenate is then added, incubated and the non-immobilized material removed, for example, by washing. A labeled species-specific antibody for the antibody sera is then added and the presence and quantity of label determined. The label determination indicates the presence of MSD in the tissue assayed. Typical

-11-

embodiments of this method include the enzyme linked immunosorbent assay (ELISA); radioimmunoassay (RIA); immunofluorescent assay (IFA); Northern, Southern, and Western Blot immunoassay.

- 5           The following examples further illustrate specific embodiments of the invention. The examples, however, are not meant to limit the scope of the invention which has been fully characterized in the foregoing disclosure.

10

#### EXAMPLE 1

- The MSD infectious agent may be characterized by determining physiochemical properties (size, sensitivity to lipid solvents, and sensitivity to  
15   protease) by treatment of the inoculum followed by the inoculation of gnotobiotic pigs to determine if the MSD infectious agent remains pathogenic.

#### A. Materials

- 20           Gnotobiotic pigs. Derivation and maintenance procedures for gnotobiotic pigs have been described in Benfield et al., Am. J. Vet. Res., 49, 330-36 (1988) and Collins et al., Am. J. Vet. Res., 50, 824-35 (1989). Sows can be obtained from a herd free of reproduction  
25   problems including MSD. Litters with stillborn and/or mummified fetuses should not be used.

- MSD inoculum (MN90-SD76-GP2, referred to herein as MN90x76-L or MN90x76-P). Trachea, lung,  
30   turbينات, tonsil, liver, brain, and spleen can be collected from nursing pigs in a Minnesota swine herd spontaneously infected with MSD (Collins et al., Minnesota Swine Conference for Veterinarians, Abstract, 254-55 (1990)). A homogenate of these tissues  
35   (designated MN 89-35477) has been prepared in Hank's Balanced Salt Solution without antibiotics and 0.5 ml can

-12-

be intranasally inoculated into three-day-old gnotobiotic piglets using a glass Nebulizer (Ted Pella Co., Redding, CA). Inoculated piglets can develop clinical signs and microscopic lesions similar to those observed in the spontaneously infected pigs. Lungs, liver, kidney, spleen, heart and brain from these gnotobiotic pigs can be collected eight days after the original inoculation and pooled to prepare another homogenate. This second homogenate can then be inoculated one additional time in gnotobiotic pigs. Again, the same tissues may be collected and homogenized, except that lung tissue can be prepared as a separate homogenate because MSD can be ideally reproduced from the lung homogenate. This lung homogenate represents the second serial passage of the original inoculum (MN 89-35477) in gnotobiotic pigs (Collins et al., 71st Meeting of the Conference of Research Workers in Animal Disease, Abstract No. 2 (1990)). Two filtrates can then be prepared using 0.20  $\mu$ m filter (Gelman Sciences, Ann Arbor, MI) and 0.10  $\mu$ m filter (Millipore Corp., Bedford, MA). These filtrates can be aliquoted and stored at  $-70^{\circ}\text{C}$ . All filtrates are free of bacteria and no viruses should be observed on direct electron microscopy using negative stained preparations.

25

Control inoculum. Homogenates of lung tissues prepared from two mock-infected gnotobiotic pigs can be used as inoculum in control pigs. This control inoculum can be prepared as 0.20 and 0.10  $\mu$ m filtrates as described for the MSD inoculum.

Necropsy procedures and histopathology. Pigs can be euthanized seven days after the original inoculation as previously described in Collins et al., 71st Meeting of the Conference of Research Workers in Animal Disease, Abstract No. 2 (1990). Tissues can be

35

-13-

collected, fixed in neutral buffered formalin, and processed for light microscopic examination as described in Collins et al., Am. J. Vet. Res., 50, 827-35 (1989). Specimens can be collected from turbinates, tonsil, 5 trachea, brain, thymus, lung (apical, cardiac, diaphragmatic lobes), heart, kidney, spleen, liver, stomach, duodenum, jejunum, ileum, ascending and descending colon, blood and mesenteric lymph nodes. These tissues can be processed and then examined using a 10 light microscope to determine whether lymphomononuclear encephalitis, interstitial pneumonia, lymphoplasmacytic rhinitis, lymphomononuclear myocarditis or portal hepatitis is present. Lesions can be consistently observed in spontaneously infected pigs from herds with 15 MSD inoculum (Collins et al., Minnesota Swine Conference for Veterinarians, Abstract, 254-55 (1990)). Fecal contents may also be collected and examined for virus particles as previously described in Ritchie et al., Arch. Gesante. Virus-forsche, 23, 292-98 (1968). Blood 20 can be collected for immunologic assays and tissues and cultured for bacteria as described in Example 3.

#### B. Infectious Agent Isolation

Lung tissue and combined brain-spleen-liver- 25 kidney tissues obtained from an infected piglet in an SIRS-infected herd were homogenized separately. Ten percent homogenates of tissue were used. The individual homogenates were mixed with Minimum Essential Medium (MEM) containing gentamicin at about 100 µg per ml. Both 30 samples were centrifuged at about 4000 x g for about 25 minutes. The supernatant was then removed and filtered through a 0.45 micron filter. The tissue and lung homogenates were then combined, and the combined material was used to infect various tissue culture cell lines.

-14-

1. In vitro testing. Two tests were conducted using 75 cm<sup>2</sup> plastic bottles. In test no. 1, the combined material was inoculated into two bottles of full cell sheet of each of the cell lines listed below.

- 5 Additionally, to one bottle of each cell line about 2.5 mg of trypsin was added. All other remaining conditions were the same for each bottle of cell line. Serum was not in the culture medium. The inoculum was 1 ml. All bottles were held for seven days at approximately 34°C.
- 10 The results were recorded at the end of seven days. After freezing and thawing, a sample was taken for a second passage in the same cell line. The remaining material was frozen and stored at about -60°C.

- In test no. 2, the combined material was
- 15 inoculated into one bottle of the same cells as were used in test no. 1. However, the cell sheets were only 20-40 percent confluent at the time of inoculation. The media contained about 10 percent fetal calf serum. Again, the inoculum was 1 ml, and the cultures were incubated at
- 20 about 34°C for approximately seven days. The results of both test no. 1 and test no. 2 are summarized below:

-15-

Cell Line Used	Test No. 1	Test No. 2
Bovine Turbinate (BT)	-	-
Feline Kidney (CRFK)	-	-
Monkey (Vero) Kidney	-	-
5 Monkey (Vero) Lung	-	-
Canine Kidney (MDCK)	-	-
Porcine (PK2a) Kidney	-	-
Mink Lung	-	-
Ferret Lung	-	-
10 Bovine Lung	-	-
Buffalo Lung	-	-
Bovine Kidney (MDBK)	-	-
Swine Testicle (ST)	-	-
Monkey Kidney (MA-104)	-	+
15 Human Rectal Tumor (HRT-18)	-	NT
Human Lung	NT	-

+ = CPE effect

- = no CPE effect

20 NT = not tested

There was no cytopathic effect observed in test no. 1 in any of the cell lines evaluated. In test no. 2, however, small clumps of MA-104 cells began to swell and form "weak holes" in the monolayer around the edges of the bottle. Fluid was separated from the bottle, passed into a new bottle of MA-104 cells (again 20-40 percent cell sheet), and then subsequently passed a third time. The cytopathic effect (CPE) became stronger with each passage. The above-described procedures were repeated for the MA-104 cell line employing a full cell sheet. CPE was also observed. Further testing demonstrated that the viral agent will also grow at 37°C. The presence of serum may be helpful for the initial isolation of the viral agent. Subsequent passages of the viral agent in the MA-104 cell line will produce the CPE without the presence of serum. However, more pronounced CPE is

-16-

observed with the use of serum in the growth medium for the MA-104 cell line.

The viral agent was passaged eight times in the MA-104 cell line with good CPE developing in three days at passage five and greater. The titer obtained is approximately 5-1/2 logs ( $10^{5.5}$ ). The viral agent will also grow in additional simian cell lines.

2. In vivo testing. A third passage harvest was used to inoculate two three-day-old gnotobiotic piglets. Both piglets were exposed intranasally, one with 1 ml and the other with 2 ml. The piglets were observed for seven days, and then were euthanized.

Tissue samples were collected for histopathologic examinations and for recovery of the viral agent. The histopathology report confirmed that lung lesions in the infected piglets were identical to lung lesions from piglets known to have SIRS. The tissue samples were processed as before, and then cultured on 20-40 percent and 100 percent monolayers of the MA-104 cell line with bovine fetal serum. The viral agent was again recovered.

A third passage harvest was also used to inoculate sows in order to verify that the reproductive effects of the disease can be duplicated and confirmed. Two multiparous sows were inoculated intranasally at 93 days of gestation. The sows delivered litters with 50 percent stillbirth piglets (8/13 and 6/14 stillborn/live) on days 112 and 114 of gestation, respectively. Seven of the stillborn piglets were partial mummies and the liveborn piglets were weak and failed to nurse vigorously. The viral agent was recovered from tissues of the stillborn piglets.

The viral agent has been recovered from three herds known to have SIRS. Antibody titers to the ATCC VR-2332 agent have been identified in these same herds.

-17-

Although there are some differences in clinical signs, i.e., cutaneous cyanosis of the ears, tail and udder in European swine, the prevailing opinion is that the North American and European diseases are caused by the same virus, a fastidious, non-hemagglutinating enveloped RNA virus as exemplified by the deposit ATCC VR-2332.

#### EXAMPLE 1A - FURTHER INFECTIOUS AGENT CHARACTERIZATION

##### 10 A. Materials and Methods

1. Cells. Crandell feline kidney (CRFK), monkey kidney (MA-104) cells were grown at 37°C in appropriate cell culture flasks. The CRFK and MA-104 cells were propagated in Eagle's minimum essential media (MEM) (available from Gibco Laboratories, Grand Island, NY) supplemented with 10 percent gamma-irradiated fetal bovine serum (FBS) (available from JRH Biosciences, Lenexa, KS), 1 percent penicillin-streptomycin and 2.5 µg/ml of amphotericin B. MA-104 cells were propagated in the same media supplemented with 10 percent FBS and 50 µg/ml of gentamicin. The FBS and cells were confirmed free of bovine virus diarrhea virus (BVDV) using previously described methods of Mayer et al., Vet. Microbiol., 16, 303-314 (1988); Smithies et al., Proc. Annu. Meet. U.S. Animal Health Assoc., 73, 539-550 (1969); and Vickers et al., J. Vet. Diagn. Invest., 2, 300-302 (1990).

2. The source of the VR-2332 isolate (SIRS virus). The source and isolation of the SIRS virus for this Example is set forth below. Virus used in this study was on the 5th to 7th passage in MA-104 cells with titers of  $10^5$  to  $10^6$  TCID<sub>50</sub>/ml.

35 Gnotobiotic pigs. Gnotobiotic piglets obtained by closed hysterotomy were maintained in stainless steel

-18-

tubs covered by flexible film isolators as previously described by Miniatas O.P. et al., Can. J. Comp. Med., 42, 428-437 (1978). The isolators were maintained at an ambient temperature of 30°C and pigs were fed recommended amounts of commercial milk substitute three times a day. Fecal swabs were collected prior to experimental inoculation and at necropsy, and were inoculated onto sheep blood agar, tergitol-seven agar and brilliant green agar in aerobic and anaerobic atmospheres. Feces collected at necropsy were also examined for viruses by negative contrast electron microscopy as described by Richie et al., Arch. Gesante. Virus-forsche, supra.

Source of Inoculum. A 160-sow farrow-to-finish herd in West Central Minnesota experienced an outbreak of MSD with typical MSD symptoms. A live sow, live neonatal piglets and stillborn fetuses were submitted to the Minnesota Veterinary Diagnostic Laboratory for examination including gross necropsy, histopathology and routine microbial investigation. An inoculum was prepared for experimental use with several tissues from clinically ill neonatal pigs. More specifically, two live and two dead 7- to 10-day-old piglets obtained during the epizootic from the affected herd were necropsied and specimens were collected for diagnostic examinations. The live piglets were euthanized by intravenous injection of euthanasia solution before necropsy. A 10 percent homogenate (MN89-35477) of brain, lung and tonsil pooled from each pig was prepared using Hank's Balanced Salt Solution (HBSS) containing 100 IU penicillin, 100 µg/ml streptomycin, and 5 µg/ml amphotericin B.

Experimental Transmission. A series of 14 gnotobiotic piglets was challenged at three days of age with pooled tissue homogenates. Each piglet was

-19-

challenged intranasally by use of a rubber bulb attached to a glass Nebulizer placed in front of the nares of the pig. Initially, two gnotobiotic piglets were inoculated with 0.5 ml each of the unfiltered inoculum (MN89-35477), 5 monitored for clinical signs of disease, and were euthanized by electrocution seven days post-exposure (PE).

A 10 percent homogenate (designated MNSD-1) of lung tissues pooled from the aforementioned gnotobiotic 10 piglets was blind passaged by exposing each of three gnotobiotic piglets to 0.5 ml of homogenate, one piglet receiving 0.5 ml of unfiltered homogenate, the second receiving 0.45  $\mu$ m filtrate, and the last one receiving a 0.22  $\mu$ m filtrate. The piglets were euthanized by eight 15 days PE and tissues were collected for histologic examination, for further passaging in gnotobiotic piglets, and for virus isolation.

A 25 percent suspension of lung (MNSD90x76-L) and a composite of brain, liver and kidney (MNSD90x76-P) 20 of the piglet inoculated with 0.45  $\mu$ m filtrate of MNSD-1 was prepared using phosphate buffered saline containing 0.5 mg/ml each of kanamycin, streptomycin, and vancomycin. Six gnotobiotic piglets were inoculated with lung homogenate MNSD90x76-L; four piglets received a 0.45 25  $\mu$ m filtrate and two were given a 0.1  $\mu$ m filtrate. Three uninfected, control gnotobiotic piglets were inoculated, one piglet with a 0.45  $\mu$ m filtrate of uninfected gnotobiotic piglet tissue homogenate in HBSS and two piglets with HBSS alone.

30

Virus Isolation. Tissue homogenates (MNSD90x76-L and MNSD90x76-P) were centrifuged at 1500 x g at 4°C for 20 minutes. The supernatant was diluted 1:1 with minimum essential medium (MEM) containing 10  $\mu$ g/ml 35 gentamicin, mixed thoroughly using a Vortex mixer and recentrifuged at 4500 x g at 4°C for 30 minutes. The

-20-

supernatant was collected and filtered through a 0.45  $\mu$ m filter. The filtrates from lung and tissue pool homogenates were combined and inoculated onto continuous cell line MA-104. Virus isolation was done in 75 cm<sup>2</sup> 5 flasks with 20-40 percent confluent monolayers of MA-104 cells containing 50 ml of MEM (pH 7.5) with 10 percent fetal bovine serum (FBS). Cell cultures were maintained at 34°C for seven days. If no cytopathic effect (CPE) was observed within seven days, cultures were frozen, 10 thawed and inoculated on MA-104 cells and incubated as above.

Virus Titration. Virus titration was done in 96-well, flat-bottom microtiter plates. Serial 10-fold 15 dilutions of virus were prepared in MEM with 2 percent FBS. After three days, the cell growth medium was drained from the microtiter plates, 200  $\mu$ l of the virus dilution was placed into each of five wells, and the plates were incubated at 37°C in an atmosphere of 5 20 percent CO<sub>2</sub>. After three days, media in wells with no CPE were replaced with MEM supplemented with 2 percent FBS (pH 7.5) and a final reading was made on the fifth day of incubation. Titers were calculated by the method of Reed et al. in Am. J. Hyg., 27, 493-497 (1938).

25

3. Other viruses. Attenuated poliovirus, available from Dr. Roger Koment, Department of Microbiology, University of South Dakota School of Medicine, Vermillion, SD, was propagated on MA-104 cells 30 to a titer of 10<sup>8</sup> TCID<sub>50</sub>/ml and the Shope strain of pseudorabies virus, available from National Veterinary Services Laboratory, Ames, IA, was grown on CRFK cells to a titer of 10<sup>5-7</sup> TCID<sub>50</sub>/ml. These viruses were used as RNA and DNA virus controls in studies to determine the 35 nucleic acid type of the VR2322 isolate of SIRS virus.

-21-

4. Preparation of antisera to VR-2332 isolate.

Passage five of the VR2322 isolate of SIRS virus (titer  $10^6$  TCID<sub>50</sub>/ml) was inactivated with 0.25 percent formalin, mixed 1:1 with Freund's incomplete adjuvant, and a rabbit was injected subcutaneously with 2 ml of this suspension at two-week intervals for six weeks. Antisera prepared two weeks after the last injection had a 1:512 neutralizing titer.

- 10                    5. Virus neutralization test (VNT). MA-104 cells were seeded in flat-bottom, 96-well microtiter plates for the VNT. Serial two-fold dilutions of each serum (100  $\mu$ l) were prepared in MEM diluent and mixed with an equal volume (100  $\mu$ l) of isolate VR-2332
- 15 containing 100-300 TCID<sub>50</sub>/100  $\mu$ l. Each mixture was incubated at 37°C for one hour, and 200  $\mu$ l of each serum-virus mixture was added to triplicate wells. Microtiter plates were incubated for an additional three days at 37°C, examined by light microscopy for cytopathic effects
- 20 (CPE), and the endpoint titer expressed as the reciprocal of the highest serum dilution which neutralized CPE.

The following polyclonal antisera, unless indicated otherwise, were prepared as described for the VR-2332 isolate (except viruses were not inactivated) and

25 used in the VNT: the Miller strain of transmissible gastroenteritis; porcine rotavirus serotype 4 (Gottfried); porcine rotavirus serotype 5 (OSU); porcine reovirus serotype 1; porcine enterovirus serotypes 1-8, available from National Veterinary Services Laboratory,

30 Ames, IA; monoclonal antibody to porcine parvovirus, available from American Type Culture Collection, Rockville, MD; encephalomyelocarditis virus, available from Dr. W. Christianson, Department of Clinical and Population Sciences, University of Minnesota, St. Paul,

35 MN; pseudorabies virus, available from National Veterinary Services Laboratory, Ames, IA; Eastern,

-22-

Western and Venezuelan equine encephalitis viruses; monoclonal antibody D89 to BVDV described by Mayer et al., Vet. Microbiol., 16, 303-314 (1988); equine arteritis virus, available from National Veterinary Services Laboratory, Ames, IA; rubella; and lactic dehydrogenase virus, available from Dr. W. Cafruny, Department of Microbiology, University of South Dakota School of Medicine, Vermillion, SD.

10           6. Direct electron microscopy (DEM). Cesium chloride gradient fractions were examined by DEM for virus particles as previously described by Benfield et al., J. Clin. Microbiol., 16, 186-190 (1982); Horzinek, "Non-arthropod-borne Togaviruses" (Acad. Press, London, 15 1981); and Richie et al., Arch. Gesante. Virus-forsche, supra, and examined on an electron microscope (Hitachi HU12A, Hitachi, Tokyo, Japan) at a potential of 75 kV.

          7. Immune electron microscopy (IEM). Immuno-  
20 gold labelling was done using goat anti-rabbit IgG gold colloidal particles (5 nm). Briefly, virus was concentrated at 40,000 x g for 30 minutes at 4°C, the pellet was resuspended in 50 µl of distilled water, and 25 µl was placed on a piece of parafilm. A collodion  
25 carbon-coated grid was floated on the drop of virus for 15 minutes, blotted dry with filter paper, and placed on a 25 µl drop of rabbit anti-SIRS sera for two minutes. Grids were washed in two changes of 0.1 percent bovine serum albumin (BSA)-Tris buffer for five minutes each and  
30 floated for 15 minutes on a drop of gold-labeled antirabbit-IgG. After six washes in BSA-Tris buffer for two minutes each, and two washes in distilled water, the grids were negatively stained and examined as described for DEM.

-23-

8. Hemagglutination test (HAT). The ability of VR-2332 isolate to hemagglutinate sheep, goat, swine, cattle, mouse, rat, rabbit, guinea pig, human type "O", duck, and chicken erythrocytes was determined using 5 standard methods. Two-fold dilutions of the 5th to 7th passage of the VR-2332 isolate of SIRS virus were prepared in phosphate-buffered saline (pH 7.2-7.4) in U-bottom microtiter plates. Equal volumes of a 1 percent suspension of washed erythrocytes from each of the above 10 species were added to each virus dilution, incubated at 4°, 22° or 37°C; and read after one to two hours when erythrocyte controls (containing no virus) had settled into a button on the bottom of the well.

15 9. Immunofluorescence (ImF). Indirect or direct ImF staining of isolate VR-2332 infected and uninfected MA-104 cells was done using the polyclonal or monoclonal antibodies tested by VNT. Swine influenza, available from National Veterinary Services Laboratory, 20 Ames, IA (type A, H1N1) and hog cholera virus polyclonal antisera, available from National Veterinary Services Laboratory, Ames, IA, were also used. Scrapings of the cell monolayer were removed at 72 hours PI with a sterile inoculating loop ImF staining as previously described by 25 Benfield et al., J. Clin. Microbiol., supra. Positive control slides were stained either with convalescent antisera from a sow (VNT titer 1:256) or a monoclonal antibody (SDOW 12 or SDOW 17, described herein) to the VR-2332 isolate of SIRS virus.

30

10. Filtration studies to estimate the size of isolate VR-2332. Clarified-infected cell culture supernatants were filtered through 0.45 µm (Schleicher and Schnell, Keene, NH), 0.20 µm (Schleicher and Schnell, 35 Keene, NH), 0.10 µm (Millipore Products Div., Bedford, MA) and 0.05 µm (Millipore Products Div., Bedford, MA)

-24-

filters. The infectivity titer before and after filtration was determined using a microtiter assay and a previously described method of Cottral, Manual of Standardized Methods for Veterinary Microbiology, pp. 81-82 (Cornell Univ. Press, Ithaca, NY, 1978). A 100-fold reduction in infectivity titer was considered significant.

#### 11. Gradient purification of isolate VR-2332.

10 The VR-2332 isolate of SIRS virus from clarified culture supernatant was concentrated by centrifugation (Beckman SW 41 Ti rotor) at 200,000 x g at 4°C for 16 hours through a discontinuous sucrose gradient consisting of 2 ml of 20, 30, 40, 50, and 65 percent sucrose (wt/vol) in 15 TNC buffer (10 mM Tris, 100 mM NaCl and 2 mM CaCl<sub>2</sub>, pH 7.8). Culture supernatants were also extracted with 1,1,2-trichlorotrifluoroethane (Sigma Chemical Co., St. Louis, MO), concentrated by ultracentrifugation at 100,000 x g at 4°C for one hour, and purified on cesium 20 chloride density gradients (1.20 g/ml) as described for the sucrose gradients. Fractions from either gradient were harvested from the top, the refractive index determined using a refractometer, and selected fractions diluted in Hank's Balanced Salt Solution for virus 25 titration as described above.

#### 12. Chloroform and fluorocarbon inactivation.

Equal volumes of SIRS virus (VR-2332) and chloroform were mixed periodically for 30 minutes at ambient room 30 temperature and then centrifuged at 500 x g at 4°C for 15 minutes. Similarly equal volumes of 1,1,2-trichlorotrifluoroethane and SIRS virus were mixed on a vortex for five minutes and centrifuged as described for the chloroform treated virus. After centrifugation, the 35 aqueous phase of each treated sample was harvested, diluted, and a microtiter assay used to determine the

-25-

remaining virus infectivity. A 100-fold reduction in titer of treated compared to untreated virus was considered significant.

- 5                    13. Effects of inhibitors of DNA viruses on  
                  isolate VR-2332. The compounds 5-bromo-2-deoxyuridine  
                  (BUDR) and mitomycin C are known inhibitors of the  
                  replication of DNA viruses, but do not inhibit RNA  
                  viruses with the exception of the Retroviridae  
10 (Easternbrook, Virology, 19, 509-520 (1962); and Reich et  
                  al., Proc. Natl. Acad. Sci., 47, 1212-1217 (1961)).  
                  Pseudorabies and poliovirus were used as the known DNA  
                  and RNA virus controls in this experiment. Cells (CRFK  
                  or MA-104) were seeded in 96-well microtiter plates and  
15 duplicate wells were inoculated with 100  $\mu$ l of ten-fold  
                  dilutions of pseudorabies virus, poliovirus or SIRS  
                  virus, respectively. After a one-hour absorption period  
                  at 37°C, media containing unabsorbed virus was removed  
                  and replaced with either MEM (untreated virus controls);  
20 MEM supplemented with 40 or 150  $\mu$ g/ml of BUDR; or MEM  
                  containing 2, 10 or 20  $\mu$ g/ml of mitomycin C. Microtiter  
                  plates were incubated at 37°C for an additional four  
                  days, examined by light microscopy for CPE, and the virus  
                  titer calculated as described previously by Cottral,  
25 supra (1978). A 1000-fold reduction in virus infectivity  
                  titer compared to the untreated control was considered  
                  significant.

14. Temperature stability of SIRS virus. A 2  
30 ml aliquot of virus in MEM was incubated either at 4°C or  
                  in water baths at 37° and 56°C for at least five days.  
                  Aliquots from each tube at the three different  
                  temperatures were collected at 15, 30, 60 and 120 minutes  
                  and then at 12-hour intervals from 12 to 120 hours.  
35 Virus samples were diluted ten-fold in MEM and virus  
                  titers calculated as described above.

-26-

## B. Results

### 1. Cytopathic effect of isolate VR-2332 on MA-

104 cells. The CPE started as small, rounded clumps of 5 cells, which appeared to be raised above the remainder of the uninfected monolayer (Figure 1B). The number of rounded cells increased and many cells became pyknotic and detached from the monolayer within two to four days PI. By five to six days PI, CPE was evident in 100 percent of the monolayer. Infectivity titers varied from  $10^5$  to  $10^7$  TCID<sub>50</sub>/1 ml on the 5th and 7th serial passage of virus, respectively. No CPE was observed in uninoculated MA-104 cells (Figure 1A).

Fluorescence was not observed in uninoculated MA-104 cells (Figure 2A). Figure 2B demonstrates the intense, diffuse fluorescence observed in the cytoplasm of inoculated MA-104 cells stained with either convalescent sow sera, rabbit antisera or monoclonal antibody (prepared to the VR-2332 isolate of SIRS).

20

### 2. Effects of lipid solvents on the infectivity of SIRS virus (isolate VR-2332).

Pretreatment of virus with chloroform eliminated infectivity (titer reduced from  $10^5$  to  $<10^1$  TCID<sub>50</sub>/1 ml), whereas fluorocarbon treatment had no significant effect.

25

### 3. Hemagglutination. Virus did not

hemagglutinate erythrocytes from 11 different species irrespective of the temperature of incubation.

30

### 4. Estimation of the size of the SIRS virus by filtration.

Virus titers were unchanged after filtration through 0.45, 0.20 and 0.10  $\mu$ m filters. However, infectivity titers were reduced 1000-fold ( $10^5$  to  $10^2$  TCID<sub>50</sub>/1 ml) after passage through a 0.05  $\mu$ m (50 nm) filter.

35

-27-

5. Effects of DNA inhibitors on replication of the SIRS virus. Only the infectivity of the DNA virus, pseudorabies, was reduced by either BUdR or mitomycin C.

5 The replication of poliovirus (RNA control) and isolate VR-2332 were not affected, indicating that the genome of the SIRS virus was probably RNA (Tables 1 and 2).

6. Virus purification. Virus bands were not  
10 detected on sucrose gradients, and peak virus titers ( $10^4$  TCID<sub>50</sub>/1 ml) were recovered from fractions with buoyant densities of 1.18 to 1.23 g/ml. This peak virus titer was 1000-fold less than the infectivity titer ( $10^7$  TCID<sub>50</sub>/ml) of the virus suspension before purification.  
15 Cesium chloride gradients yielded a single, faint, opalescent band at a buoyant density of 1.18 to 1.19 g/ml. Peak virus infectivity of  $1.3 \times 10^6$  TCID<sub>50</sub>/1 ml, which was 130 times higher than peak infectivity from the sucrose gradient, corresponded to this visible band  
20 (Figure 3).

7. DEM of purified SIRS virus particles.  
Pleomorphic, but predominantly spherical virions were observed by DEM in the CsCl fractions (1.18 to 1.19  
25 g/ml). Virions were 48 to 80 nm (average of 25 particles = 62 nm) in diameter and consisted of complete, electron translucent or empty (electron dense center) particles surrounded by a thin membrane or envelope (Figure 4A). [A 40 to 45 nm diameter isosahedral nucleocapsid with  
30 short surface projections of about 5 nm in length was observed.] These particles contained cores that were 25 to 35 nm in diameter (Figure 4B). Virions were immuno-gold labeled if the rabbit hyperimmune antisera and gold-labeled anti-rabbit IgG were used as primary and  
35 secondary antibodies (Figure 4B). Virions were not gold-

-28-

labeled in the absence of the rabbit hyperimmune antisera to isolate VR-2332.

8. Antigenic relationship of SIRS virus to  
5 antisera from other viruses. Antisera to known porcine  
viruses and various togaviruses did not neutralize or  
react with SIRS virus antigen in infected MA-104 cells.  
Convalescent sow and hyperimmune rabbit antisera had  
neutralizing antibody titers of 1:256 and 1:512,  
10 respectively.

9. Effects of temperature on the infectivity  
of the SIRS virus. Virus infectivity for MA-104 cells  
was reduced 50 percent after incubation for 12 hours at  
15 37°C and completely inactivated after 48 hours of  
incubation at 37°C and 45 minutes incubation at 56°C  
(Figure 5). Infectivity was unchanged after one month  
incubation at 4°C or four months at -70°C (data not  
shown). Lung tissue collected from gnotobiotic pigs  
20 infected with SIRS virus and homogenized in Hank's  
Balanced Salt Solution retains infectivity for pigs for  
at least 18 months.

Results indicate that isolate VR-2332 is a  
fastidious, non-hemagglutinating enveloped RNA virus,  
25 which can be tentatively classified as a non-arthropod  
borne togavirus belonging to an unknown genera.

The presence of an RNA genome of the SIRS virus  
was confirmed by the ability of this virus to continue to  
replicate in the presence of 5-bromo-2-deoxyuridine and  
30 mitomycin C, which are known to inhibit the replication  
of DNA and one family of RNA viruses (Retroviridae)  
(Easterbrook, supra (1962); and Reich et al., supra  
(1961)), but not other RNA viruses. Our provisional  
classification of the SIRS virus as an RNA virus agrees  
35 with the observation that this virus replicates in the

-29-

cytoplasm of the cell as indicated by the presence of virus antigens detected by ImF.

The VR-2332 isolate of SIRS virus is heat labile, but relatively stable for long periods of time at 4° and -70°C. The thermolability of this virus at 37°C has practical applications for propagation of the virus, suggesting that growth at temperatures lower than 37°C will produce higher virus yields. Refrigeration is sufficient for preservation of diagnostic specimens for virus isolation for short periods of time, otherwise the sample can be frozen for several months or longer.

#### EXAMPLE 2

The purest form of an inoculum with the MSD infectious agent as determined from experiments in Example 1 may be used to transmit the MSD agent to pregnant sows to reproduce the reproductive form of the disease syndrome.

Discussion. Recently, transient anorexia and premature farrowing (both prominent clinical signs of MSD in the field) was induced in 2/2 sows inoculated with the same MSD inoculum, which produces respiratory lesions in gnotobiotic pigs. In addition, 15/29 (52 percent) of the pigs were stillborns and the remaining 14 pigs were weak and did not nurse well. No gross or microscopic lesions were observed in the stillborn pigs or the placenta and isolation procedures to detect microbial agents are now in progress. Therefore, the experiment described below tests whether the reproductive form of MSD can be transmitted to sows by intranasal inoculation and whether interference with fetal viability results from replication in maternal tissues but not fetuses.

Experiments in Example 1 provide information on how the inoculum can be treated (filter size, organic solvent extraction and/or protease digestion) to provide

-30-

the purest form of the MSD agent for an inoculum (i.e., viral agent). Because the MSD agent has both a respiratory form in young pigs and a reproductive form in adult swine, it is necessary to reproduce the latter form of the syndrome to further verify that the infectious agent is the putative cause of MSD.

A 93-day gestational sow is used as the experimental animal because it is possible to experimentally induce abortion in these animals inoculated with the MSD infectious agent. Sows can be purchased from a commercial herd free of ongoing reproductive problems and MSD. Complete epidemiologic records on this herd can be computerized and information on gestation times, litter sizes, and average number of stillbirths can be made available for comparative studies. Groups of three sows each can be intranasally inoculated at 93 days of gestation with either the 0.20  $\mu$ m filtrate (positive controls), a pathogenic but modified inoculum as dictated by results from Example 1, a 0.20  $\mu$ m filtrate of the control inoculum (negative control), and a control inoculum modified as indicated by results of experiments in Example 1. Each group of sows can be housed in separate isolation rooms and examined daily until gestation is complete. Temperatures and clinical signs (anorexia, respiratory problems such as coughing, sneezing, panting, and increased respiration) can be noted daily. Sampling of sows can be restricted to a pre- and post-farrowing blood sample for serology. The actual date of farrowing can be noted and the number of stillborns, mummified fetuses, live "weak" pigs and live "normal" pigs determined. Fetuses can be examined for gross and microscopic lesions as described in Example 1 and fetal tissues processed for microbiologic assays as described in Example 3. The fetal sera can also be assayed for the presence of gammaglobulins and antibodies to PPV and EMCV (Joo et al., In Proceedings of the

-31-

Mystery Swine Disease Committee Meeting, 62-66 (1990);  
and Kim et al., J. Vet. Diagn. Invest., 1, 101-4 (1990)).  
Pigs born live can be observed for one week and morbidity  
and mortality recorded, after which these pigs can be  
5 euthanized and the tissues collected for light  
microscopic and microbiologic examination as described  
for the fetuses.

The 0.2  $\mu$ m and the modified filtrates of the  
MSD inoculum are pathogenic for sows and induce anorexia,  
10 possibly a mild fever, and premature farrowing with a  
large number of stillborn and weak pigs in each litter.  
This illustrates evidence that the inoculum contains the  
MSD agent. Sows inoculated with the control inoculum  
farrow near term and have litter sizes within the normal  
15 range for the herd of origin as determined from the  
available epidemiologic database on this herd. No  
lesions in the stillborn pigs are found and a high rate  
of mortality among the surviving weak pigs within one  
week after birth is observed.

20

### EXAMPLE 3

Tissue samples collected from gnotobiotic  
piglets inoculated with the MSD inoculum and euthanized  
at various times post-inoculation can be used to isolate  
25 and identify the MSD infectious agent to determine the  
sequential development of lesions, and to ascertain  
whether the MSD agent is immunosuppressive.

Immunofluorescence assays on frozen tissues and  
30 inoculated cell cultures. Immunofluorescence assays on  
frozen tissues and cell scrapings can be done as  
previously described in Benfield et al., J. Clin.  
Microbiol., 16, 186-190 (1982) and Benfield et al., Am.  
J. Vet. Res., 49, 330-36 (1988). The frozen tissues and  
35 cell scrapings can be screened for PPV, EMCV (See Example  
4 for definition of acronyms) and the MSD agent(s).

-32-

Conjugates for PPV and EMCV are available at the South Dakota Animal Disease Research and Diagnostic Laboratory. A hyperimmune sera in gnotobiotic pigs can be prepared from the purest form of the MSD inoculum. Two pigs can be inoculated intranasally, and then given a subcutaneous booster of the MSD inoculum in Freund's incomplete adjuvant at two and four weeks after the initial inoculation. Sera can be harvested from this pig two weeks after the last booster. A control sera is also prepared in two gnotobiotic pigs using the control inoculum and the same immunization protocol described for the MSD inoculum. These sera can be used as primary antibody and goat or rabbit anti-porcine immunoglobulin conjugated with fluorescein isothiocyanate as secondary antibody to detect MSD antigens in frozen tissue sections and cell cultures.

Serologic assays. Sera collected from control and inoculated pigs can be assayed for the presence or absence of antibody to PPV and SIV (hemagglutination inhibition), Leptospira (micro-agglutination), and EMCV (viral neutralization) (see Example 4 for definition of acronyms). Previous results indicated that serology to other common microbial agents were negative (Collins, et al., 71st Meeting of the Conference of Research Workers in Animal Disease, Abstract No. 2 (1990)).

Immunologic assays. Tissues and blood can be collected from MSD inoculated and control pigs so that their immunological status can be determined. Porcine leukocytes can be isolated from peripheral blood by single step discontinuous gradient floatation on Histopaque 1077 as taught by Pescovitz et al., J. Immunol., 134, 37-44 (1985). Cells from lymph nodes, spleen and thymus can be spilled into single cells by moderate mincing of the tissues and collection of the

-33-

resultant cell suspensions (Hurley et al., Cancer Res., 47, 3729-35 (1987)).

Porcine leukocyte phenotypes can be determined using a panel of monoclonal antibodies available through the American Type Culture Collection and Joan Lunney (USDA Beltsville, MD). These include antibodies for the measurement of total T cells, pCD2 (MSA4; Hammerberg et al., Vet. Immunol. Immunopathol., 11, 107-21 (1986), helper/class II MHC dependent T cells, pCD4 (74-12-4; Lunney et al., Vet. Immunol. Immunopathol., 17, 135-144 (1987), cytotoxic-suppressor/class I MHC dependent T cells, pCD8 (74-2-11; Ibid), macrophages and granulocytes (74-22-15A; Ibid), thymocytes and peripheral B cells, pCD1 (76-7-4; Ibid), and pig MHC class II antigens equivalent to human DRw (MSA3; Hammerberg et al., Vet. Immunol. Immunopathol., 11, 107-21 (1986)) and DQw (TH21A and others (VRMD, Pullman, WA; Davis et al., Hybridoma Technology in Agriculture and Veterinary Research, Rowman and Allanheld, 121-50 (1984)). Isotype-specific monoclonal antibodies to porcine immunoglobulins are also available (Paul et al., J. Vet. Res., 50, 471-79 (1989)), and can be used at twice minimum saturating concentration for indirect fluorescent staining of leukocytes from peripheral blood, lymph node, and Peyer's patches (Hurley et al., Vet. Immunol. Immunopathol., 25, 177-93 (1990)). To achieve two-color analysis, cells can also be stained with rPE-labeled avidin after being tagged with biotin-bound (Pierce kit #21333) antibodies. Cells can be analyzed by flow cytometry or a two-color analytical fluorescent microscope (PTI FSCAN system). Co-detection in the 488 nm laser line on the flow cytometer or using the dual analytical fluorescent microscope can easily be attained. Intensity of cellular fluorescence and percentage of positive cells can also be determined.

In vitro functional assays such as lectin mitogenesis with concanavalin A, pokeweed mitogen (PWM),

-34-

and phytohaemagglutinin (PHA), can be performed as described in Hammerberg et al., Am. J. Vet. Res., 50, 868-74 (1989). Antigen-specific in vitro T cell responses to lysozyme can also be modeled after their technique. B cell proliferative assays can be performed with E. coli and S. typhimurium LPS or anti-immunoglobulin as reported in Symons et al., Int. Archs. Allergy Appl. Immun., 54, 67-77 (1977). In vitro antibody production, induced with PWM, can be accomplished and quantitated as described in Hammerberg et al., Am. J. Vet. Res., 50, 868-74 (1989). Macrophage production of IL-1 after 48-hour exposure to E. coli LPS can be measured in the mouse thymocyte assay (Mizel, Immunological Rev., 63, 51-72 (1982)). IL-2 production by PHA-stimulated lymphocytes can be measured as described by Stott et al., Vet. Immunol. Immunopathol., 13, 31-38 (1986). An isotype-specific anti-lysozyme ELISA can be done utilizing the monoclonal antibodies to porcine immunoglobulin isotypes (Paul et al., Am. J. Vet. Res., 50, 471-79 (1989)).

To assess in vivo antibody production, three piglets inoculated with MSD and three inoculated with control inoculum can be injected with a 2 percent suspension of sheep erythrocytes and a 10 µg/ml solution of bovine serum albumin at separate sites at 5, 7, 10, 14, and 24 days after their original inoculation. Pigs are euthanized at 24 days after the original inoculation, tissues are collected for histopathology as described in Example 1, and blood is collected to assay for antibody.

The total antibody level and the specific IgG and IgM responses to each antigen can be measured by antigen-specific radial immunodiffusion or ELISA. Antigen-specific plaque assays can be performed on spleen cells to assess the frequency of B cell clones in the infected and control animals (Kappler, J. Immunol., 112, 1271-85 (1974)).

-35-

EXAMPLE 4

Objective. The goal of this experiment is to prepare hyperimmune antisera in a gnotobiotic pig to an isolate of MSD for use as a diagnostic reagent and for further characterization of the antigenic properties of MSD.

Background. Previous studies done in gnotobiotic pigs at South Dakota State University in collaboration with the University of Minnesota indicated that pooled tissue homogenates from field case MN 89-35477 induced lung lesions in gnotobiotic pigs. Pooled tissue homogenates from these pigs have subsequently been used to produce clinical disease and respiratory lesions characteristic of MSD in three-day-old gnotobiotic pigs (Collins et al., 71st Meeting of the Conference of Research Workers in Animal Disease, Abstract No. 2 (1990) and Collins et al., Minnesota Swine Conference for Veterinarians, Abstract, 254-55 (1990)). Lung and tissue homogenates were prepared from this second passage of the original inocula in gnotobiotic pig (90 X 75) to produce a second inocula. The second inocula can be used as inocula and antigen to produce the hyperimmune sera in this experiment.

Procedure to Accomplish the Objective

Gnotobiotic pigs. Gnotobiotic pigs can be derived and maintained as previously described in Benfield et al., Am. J. Vet. Res., 49, 330-36 (1988) and Collins et al., Am. J. Vet. Res., 50, 827-835 (1989).

-36-

Inoculation of hyperimmune sera. Hyperimmune sera can be prepared by initially inoculating one gnotobiotic pig as described above. Pigs can then be given a booster consisting of 1 ml of the second inocula and 1 ml of Freund's Incomplete Adjuvant at 14 and 21 days after the original inoculation (Harlow and Lane, 1988). The pig should then be killed and exsanguinated 14 days or later after the last inoculation. Sera should be harvested and dispensed into appropriate aliquots and frozen at -20°C.

Serology. The hyperimmune sera can be tested for the presence of antibodies to common pathogens of swine as commonly done in most diagnostic laboratories. This sera can be tested for antibodies to Hemophilus, Brucellosis, Leptospira (6 serovars), pseudorabies (PRV), parvovirus (PPV), encephalomyocarditis virus (EMC), and Swine Influenza Virus (SI).

Results. The pig used for preparation of the antisera was pig #4B (experimental number 90 X 238). This pig was inoculated on 11/1/90 and observed daily for clinical signs until killed on 11/29/90. Clinical signs are summarized in Table 1. Unfortunately the continuing degenerate condition of the pig mandated that it be euthanized after only one booster on 11/13/90. The pig was euthanized 16 days after the initial booster on 11/29/90.

Serology results were negative for all the above agents except PPV, which had a titer of 16, 384 (See Table 2). A pretitered sera on this pig was not conducted.

Samples of lung, heart, brain, kidney, colon, small intestine, turbinates, spleen, stomach and trachea were collected when the pig was necropsied. These samples were evaluated and it was found that the lungs

-37-

from this pig had lesions of severe pneumonia typical of that seen with field cases of MSD.

The results of this experiment confirm initial studies that an infectious agent is present in the second 5 inocula because it can induce clinical disease and lesions typical of those observed in natural cases of MSD.

-38-

Table 1

<u>Date</u>	<u>Observations of Inoculated Piglet (90x238)</u>
10/29	Surgery
11/1	Pig inoculated i.n. with 0.5 ml of above inocula using Nebulizer at 5:00 p.m.
11/2	Not observed.
11/3	This pig has 2 times as much milk in bowl as the control pigs. I'm not sure how well the pigs are eating or if the agent is the cause of the anorexia. Feces normal.
11/4	May be a little slow but alert and strong, 1/2 milk left, feces soft and brown.
11/5	9 a.m.: strong, alert, drank most of milk, feces mucoid and brown (2).  4 p.m.: strong, alert, drank most of milk, feces mucoid and brown (2).
11/6	9 a.m.: strong, alert, drank 1/2 of milk, feces brown mucoid (2).  6 p.m.: strong, alert, 1/2 of milk left, feces loose yellow brown (2).
11/7	9 a.m.: strong, alert, drank most of milk, feces light brown mucoid (2).  3 p.m.: strong, alert, drank most of milk, feces light brown mucoid (2).
11/8	2 p.m.: Alert, does not drink milk as well as controls, slower than controls, mucoid yellow feces.
11/9	8 a.m.: Alert, still does not drink as aggressively as controls, pasty feces.  5 p.m.: same observation as 8 a.m.
11/10	6 p.m.: alert, vigorous, rubs snout aggressively against feed pan, feces brown loose, not eating like controls.

-39-

Table 1 Cont'd

11/11	6 p.m.: alert, vigorous, rubs snout vigorously against feed pan, feces pasty, not eating as well as controls, still milk in pan, by 6:30 p.m. controls had finished eating.
11/12	9 a.m.: alert, but not as aggressive as controls, after 5 minutes the controls have cleaned pans Xut this pig still has at least 2/3 of milk in bowl. Some snout rubbing.
11/13	Inoculated with 90 X 75 lg & pool using nebulizer (.5 m) and sp. IFA (1 ml) at 4 p.m. Alert but not as aggressive as controls, still has 1/2 pan of milk but controls have consumed all their milk. No snout rubbing, feces pasty.
11/15	7 p.m.: alert but not as aggressive as controls, still eat slow, rough hair coat, but gaining weight like controls.
11/16 to 11/24	Not much change, alert but steady declining in activity  1 p.m. (11/24): respiration seems to be more rapid, hair coat is rough, not gaining weight like controls.
11/29	Euthanized - blood collected for H.I. sera. Usual times collected for histopathology including tonsil. Blood collected for lymphocyte mitogenic assays. No gross lesions noted. Set up turbinate explant cultures.

Table 2Antibody Tests of Inoculated Piglet

- |    |    |                                       |
|----|----|---------------------------------------|
| 5  | 1. | Swine influenza - negative            |
|    | 2. | Swine Encephalomyocarditis - negative |
|    | 3. | Swine APP - negative                  |
| 10 | 4. | Swine PRV - negative                  |
|    | 5. | Swine PPV - negative                  |
| 15 | 6. | Swine Brucella - negative             |
|    | 7. | Swine Leptospirose - negative         |
|    | 8. | Swine EPI - negative                  |
| 20 |    |                                       |

EXAMPLE 4A - MONOCLONAL ANTIBODY PREPARATION

Mouse immunization. Eight weeks before the date of hybridoma fusion, inoculate BALB/c AnN mice IP with a 1:1 suspension of antigen in Freund's complete adjuvant (CFA). The amount of antigen used is dependent on the immunogenicity and toxicity of the antigen. Use a maximum of 0.3 ml CFA per mouse. Five weeks after the initial immunization, boost mice with antigen in CFA.

One week prior to the fusion, immunize mice IP with antigen in saline. Two days before the fusion, inoculate mice IV with antigen in saline.

Myeloma cells. Maintain mouse myeloma cells (P3/NS-1/1-Ag4-1 (ATCC TIB 18)) on Dulbecco's Modified Eagle Media (DMEM) with 10 percent fetal bovine serum. Approximately  $10^7$  to  $10^8$  cells are required for fusion with B cells obtained from one typical mouse spleen. Immediately prior to hybridoma fusion, harvest myeloma cells from a log phase culture into a 50 ml centrifuge tube and pellet cells by centrifugation at 200 x g for five minutes. Remove the supernatant and wash cells

-41-

twice with serum free DMEM followed by centrifugation as above. Resuspend the pellet (containing  $10^7$  to  $10^8$  cells) in 1 ml serum free DMEM.

5                    Isolation of Spleen Lymphocytes

Euthanize the mouse by cervical dislocation and immerse in 70 percent ethanol for several minutes. Remove the spleen by aseptic procedure and transfer it to a sterile petri dish containing 5 ml cold serum free  
10 DMEM. Use a scalpel blade to slit the spleen along the long axis and gently scrape along the length of the spleen to release the splenocytes into the media. Use a pipette to transfer the free cells and media to a 15 ml centrifuge tube, leaving the spleen casing behind. Allow  
15 the tissue debris in the tube to settle for five minutes and transfer the single cell suspension to a fresh tube. Centrifuge cells for five minutes at 200 x g, discard the supernatant and wash the pellet again with cold serum free DMEM. Resuspend cells in 1 ml serum free DMEM and  
20 store cells on ice until fusion.

Fusion. Add the spleen cells to the centrifuge tube containing the myeloma cells and centrifuge for five minutes at 500 x g. Remove all the supernatant and  
25 loosen the cell pellet by tapping on the side of the tube. Keeping all reagents and cells at 37°C, add 1 ml 50 percent polyethylene glycol solution (PEG 4000, Gibco, Grand Island, NY) dropwise to the tube over a one-minute period with gentle mixing. Allow the mixture to stand at  
30 37°C for one minute. Add 1 ml warm serum free DMEM dropwise over one minute with gentle mixing. Finally, add 20 ml serum-free DMEM dropwise over four minutes, then immediately centrifuge cells at 200 x g for five minutes. Discard the supernatant and resuspend cells in  
35 47 ml DMEM containing 20 percent fetal bovine serum, 0.2 units/ml insulin, 0.5 mM sodium pyruvate, 1 mM

-42-

oxaloacetic acid, 2 mM L-glutamine, non-essential amino acids, and 10 percent NCTC-109 lymphocyte media. Add 1 ml volume of cell suspension to the wells of two 24-well, flat-bottom tissue culture plates. Include a myeloma cell control well and incubate plates at 37°C and 10 percent CO<sub>2</sub> (SDMEM).

Following overnight incubation, remove 0.5 ml of media from each well without disturbing the cell layer. Add 1 ml SDMEM containing  $1 \times 10^{-4}$  M hypoxanthine,  $4 \times 10^{-7}$  M aminopterin, and  $1.6 \times 10^{-5}$  M thymidine (HAT) to each well and continue incubation at 37°C and 10 percent CO<sub>2</sub>. Continue to replace 1 ml of used media with 1 ml of fresh DMEM + HAT three times weekly for two to three weeks. When significant clone growth is apparent, assay the wells for the presence of specific antibody by ELISA, indirect FA, or other appropriate assay systems.

**Cloning.** Primary wells testing positive for specific antibody should be subcloned immediately to obtain a stable cell line and avoid overgrowth by other clones. Resuspend cells from selected primary wells and perform cell counts using trypan blue stain and a hemocytometer. Make dilutions of the cells to obtain a final concentration of about 2 cells/ml in SDMDM + HT. Use normal spleen cells obtained from non-inoculated mice as a feeder layer by adding 50  $\mu$ l packed cells per 100 ml media.

Add 250  $\mu$ l of cell suspension to each well of 96-well plates and incubate at 37°C and 10 percent CO<sub>2</sub>. Clones should be visible in two to three weeks and supernatants from wells containing single clones are assayed when significant growth is apparent. Repeat the cloning procedure with wells testing positive for specific antibody. Slowly expand selected clones to tissue culture flasks for further characterization and cryopreservation.

Ascites Production. Prime BALB/c AnN mice with 0.5 ml pristane given IP two weeks before inoculation with hybridoma cells. Harvest hybridoma cells and wash 5 once with Hank's Balanced Salt Solution (HBSS). Resuspend cells in HBSS and inoculate primed mice IP with  $10^4$  to  $10^6$  viable cells. When ascites production is apparent (usually one to two weeks after inoculation) drain by inserting a 16 G 1-1/2" needle ventrally in the 10 inguinal region. Hold the hub of the needle over a centrifuge tube and drain ascites into the tube. Centrifuge ascites fluid at 200 x g, filter through a 0.2 nm filter, and store frozen.

15 EXAMPLE 4B - SIRS MONOCLONAL ANTIBODIES  
(SDOW 12 AND SDOW 17)

The SDOW 12 (ATCC No. HB 10996) and SDOW 17 (ATCC No. HB 10997) monoclonal antibodies to the SIRS virus were prepared using the above standard monoclonal antibody production protocol. The antigen used for mouse immunization was passage 6 SIRS virus (VR-2332) grown on MA-104 cells obtained from Boehringer Ingelheim Animal Health (BIAH). For each immunization, mice were inoculated with 0.3 ml of gradient purified virus with a titer of  $10^6$  TCID<sub>50</sub>/100  $\mu$ l.

An indirect fluorescent antibody (IFA) assay was used to detect specific antibody in hybridoma primary wells and clone wells. Acetone fixed virus infected and non-infected cell monolayers in 96-well tissue culture plates were used for the IFA. Cell culture supernatants and ascites fluids from these hybridomas produced bright, granular cytoplasmic fluorescence in SIRS infected cells.

Preliminary characterization of these monoclonal antibodies included immunoglobulin isotyping and radioimmunoprecipitation of SIRS viral proteins. The SDOW 12 and SDOW 17 monoclonal antibodies are both of the

IgG<sub>1</sub> isotype. They also both bind to a 15 kD viral protein on radioimmunoprecipitation.

#### EXAMPLE 5

5           Three pilot studies are described. A gnotobiotic pig study was undertaken to show that field material could be used to infect and cause the respiratory component of the syndrome in germ-free pigs. A second study using conventional weaned pigs was  
10 undertaken to determine if the respiratory disease seen in gnotobiotic pigs could be reproduced in conventional pigs. And finally, a pregnant sow study was undertaken to determine if the reproductive failure component of the syndrome could be experimentally reproduced.

15

#### Material and Methods

Field case. See Source of Inoculum in Example 1A.

20           Gnotobiotic study. Six hysterectomy-derived gnotobiotic piglets were inoculated intranasally at three days of age with the field inoculum (10 percent homogenates, various tissues). Filtered (0.22  $\mu$ m) and unfiltered inoculum were used. Two control piglets were  
25 inoculated with media only. Clinical signs were monitored daily and the pigs were euthanized eight days post-inoculation, except one animal which was held for the production of hyperimmune serum. Tissues for virus isolation and histologic examination were collected at  
30 necropsy along with sera which was screened for antibodies to leptospira, chlamydia, eperythrozoon, Aujeszky's disease virus, porcine parvovirus, encephalomyocarditis virus, hemagglutinating encephalitis virus, swine influenza virus, bovine respiratory  
35 syncytial virus, canine distemper virus, bovine viral diarrhea and hog cholera. The original inoculum and

-45-

tissues from gnotobiotic piglets were inoculated onto continuous and primary cell lines for three passages. In addition, direct and immunoelectron microscopy was performed.

5

Conventional pig study. Three conventional 28-day-old weaned pigs from a farm with no history of MSD were intranasally inoculated with 10 percent lung homogenates from affected gnotobiotic piglets. A lung homogenate from a negative gnotobiotic pig was used as inoculum for a control. Piglets were monitored daily for clinical signs and were necropsied eight days post-inoculation. Sera/tissues were processed as described above.

15

Pregnant sow study. Eight multiparous sows with known historical due dates from a farm free of MSD were used in this study. Three weeks prior to the expected farrowing date, six sows were intranasally inoculated with affected gnotobiotic lung homogenates and two sows with negative lung homogenates. Clinical signs were monitored daily. The sows were allowed to farrow naturally and when possible the farrowings were attended in order to collect presuckled sera from live born pigs. Sows and live pigs were euthanized shortly after farrowing and tissues were collected for histopathology and virus isolation. Sera or fetal thoracic fluids were also collected.

### 30 Results

Field study. No gross lesions were seen at necropsy. Microscopic examination of nursing piglets revealed necrotizing interstitial pneumonia and lymphomononuclear encephalitis. The fetuses did not have lesions but the sow did have a mild encephalitis.

-46-

Microbiologic examination did not yield conclusive results.

Gnotobiotic study. The piglets became anorexic and developed rough hair coats three days post-inoculation. Controls remained normal. Microscopic lesions were found in the principals inoculated with filtered or non-filtered material. The lesions were similar to field cases and included: necrotizing interstitial pneumonia (6/6 [six out of six piglets]), lymphoplasmacytic rhinitis (4/6), lymphomononuclear encephalitis (2/6) and myocarditis (1/6). No etiologic agent was identified either by pre- and post-inoculation serology or through inoculum/tissue examination.

15

Conventional weaned pig study. Clinically, the principals became dull and anorexic two days post inoculation. The animals appeared chilled even though adequate heat was provided. One pig had an elevated temperature (41.5°C) six days post-inoculation. Interstitial pneumonia, encephalitis and myocarditis were found in the principals but not the control.

Pregnant sow study. Clinically, only two principals showed any significant temperature rises (1.5°C) day three or five post-inoculation. However, anorexia was noted in 4/6 sows at day four or five post-inoculation. Three sows farrowed up to seven days early and three sows farrowed on time. Over 50 percent of the fetuses from infected sows were born dead while the controls had normal litters. Both stillborns and late-term mummies were found in infected litters. Laboratory findings were not conclusive--no specific agent has been identified and no lesions have been noted in fetuses to date.

-47-

Although no causative microorganism has been identified, the findings suggest MSD can be transmitted experimentally to gnotobiotic and conventional pigs (using field tissues from one farm). Both the 5 respiratory and reproductive forms of MSD were reproduced. The agent involved appears infectious, filterable at 0.22  $\mu$ m and is seemingly fastidious.

#### Discussion

10 MSD is an important emerging disease not only in the United States but also throughout the world. In order to study the disease in a controlled setting, gnotobiotic pigs were inoculated intranasally at three 15 days of age with tissue homogenates from a farm experiencing the clinical signs of MSD. Microscopic lesions similar to field cases including necrotizing interstitial pneumonia and to a lesser extent lymphoplasmacytic rhinitis, lymphomononuclear 20 not controls. Using lung homogenates from the gnotobiotic pigs, intranasal inoculation of conventional four-week-old weaned pigs produced similar lesions. Multiparous pregnant sows were also inoculated with 25 due dates. Clinically these sows went through a period of anorexia and farrowed up to seven days prior to their beginning stages of mummification. The findings indicated the disease associate with the infectious agent 30 can be isolated and transmitted experimentally with field tissues to gnotobiotic pigs and from gnotobiotic pigs to conventional weaned pigs or pregnant sows. This study provides a model for both the respiratory and 35 reproductive forms of the disease which will lead to further investigations of the pathogenesis and diagnosis of MSD.

International Application No: PCT/

**MICROORGANISMS**

Optional Sheet in connection with the microorganism referred to on page \_\_\_\_\_, line \_\_\_\_\_ of the description 1

**A. IDENTIFICATION OF DEPOSIT :**Further deposits are identified on an additional sheet ☒ (antibody) SDOW 12

Name of depositary institution \*

AMERICAN TYPE CULTURE COLLECTION

Address of depositary institution (including postal code and country) \*

12301 Parklawn Drive  
Rockville, Maryland 20852  
United States of America

Date of deposit \*

27 March 1992

Accession Number \*

HB 10996

**B. ADDITIONAL INDICATIONS** \* (leave blank if not applicable). This information is continued on a separate attached sheet: ☐**C. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE** \* (if the indications are not for all designated States)**D. SEPARATE FURNISHING OF INDICATIONS** \* (leave blank if not applicable)

The indications listed below will be submitted to the International Bureau later \* (Specify the general nature of the indications e.g., "Accession Number of Deposit")

**E.** ☒ This sheet was received with the international application when filed (to be checked by the receiving Office)Virginia L. Lily  
(Authorized Officer)☐ The date of receipt (from the applicant) by the International Bureau is:

was

(Authorized Officer)

(January 1985)

International Application No: PCT/

/

**MICROORGANISMS**Optional Sheet in connection with the microorganism referred to on page \_\_\_\_\_, line \_\_\_\_\_ of the description <sup>1</sup>**A. IDENTIFICATION OF DEPOSIT <sup>1</sup>**Further deposits are identified on an additional sheet ☒ <sup>2</sup> (antibody) SDOW 17Name of depository institution <sup>3</sup>

AMERICAN TYPE CULTURE COLLECTION

Address of depository institution (including postal code and country) <sup>4</sup>12301 Parklawn Drive  
Rockville, Maryland 20852  
United States of AmericaDate of deposit <sup>5</sup>

27 March 1992

Accession Number <sup>6</sup>

HB 10997

**B. ADDITIONAL INDICATIONS <sup>7</sup>** (leave blank if not applicable). This information is continued on a separate attached sheet ☐**C. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE <sup>8</sup>** (if the indications are not for all designated States)**D. SEPARATE FURNISHING OF INDICATIONS <sup>9</sup>** (leave blank if not applicable)The indications listed below will be submitted to the International Bureau later <sup>9</sup> (Specify the general nature of the indications e.g., "Accession Number of Deposit")**E. ☒** This sheet was received with the international application when filed (to be checked by the receiving Office)

(Authorized Officer)

*Virginia L. Lely*☐ The date of receipt (from the applicant) by the International Bureau <sup>10</sup>

was

(Authorized Officer)

International Application No: PCT/

/

## MICROORGANISMS

Optional Sheet in connection with the microorganism referred to on page \_\_\_\_\_, line \_\_\_\_\_ of the description 1

## A. IDENTIFICATION OF DEPOSIT 1

Further deposits are identified on an additional sheet ☐ (SIRS virus) BIAH-001

Name of depository institution \*

AMERICAN TYPE CULTURE COLLECTION

Address of depository institution (including postal code and country) \*

12301 Parklawn Drive  
Rockville, Maryland 20852  
United States of America

Date of deposit \*

18 July 1991

Accession Number \*

VR 2332

B. ADDITIONAL INDICATIONS 1 (leave blank if not applicable). This information is continued on a separate attached sheet ☐

C. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE 1 (If the indications are not for all designated States)

D. SEPARATE FURNISHING OF INDICATIONS 1 (leave blank if not applicable)

The indications listed below will be submitted to the International Bureau later 1 (Specify the general nature of the indications e.g., "Accession Number of Deposit")

E. ☒ This sheet was received with the international application when filed (to be checked by the receiving Office)

(Authorized Officer)

*Virginia L. Lilly*☐ The date of receipt (from the applicant) by the International Bureau is \*

was

(Authorized Officer)

(January 1985)

-51-

**WHAT IS CLAIMED IS:**

1. A vaccine suitable for use in prevention of mystery swine disease, comprising:

an inactivated or attenuated infectious agent derived from swine tissue infected with mystery swine disease in combination with a pharmaceutical carrier.

2. A vaccine according to claim 1 wherein the infectious agent is obtained from an inoculum of a filtered homogenate of swine lung tissue infected with mystery swine disease.

3. A vaccine according to claim 2 wherein the homogenate has been purified by neutralization with antibody sera to swine diseases selected from the group consisting of hemophilus, brucellosis, leptospire, parovirus, pseudorabies, encephalomyocarditis, enterovirus, swine influenza and any combination thereof.

4. A vaccine according to claim 1 wherein the infectious agent is obtained from the cell culture medium of in vitro cultured, avian, insect, or mammalian organ cells, primary or continuous, treated with an infectious agent isolated from the swine lung tissue known to have swine infertility and respiratory syndrome disease.

5. A vaccine according to claim 2 wherein the filtered homogenate contains biological particles having a size of no greater than 1.0 micron.

6. A vaccine according to claim 5 wherein the size is no greater than 0.5 micron.

7. A vaccine according to claim 1 wherein the infectious agent is a virus.

8. A vaccine according to claim 7 wherein said virus is a fastidious, non-hemagglutinating enveloped RNA virus.

9. A vaccine according to claim 1 wherein the infectious agent is a biologically pure culture of swine infertility and respiratory syndrome virus ATCC VR-2332 and zoopathogenic mutants thereof.

10. A vaccine according to claim 7 wherein the virus has been purified by gradient or serial cell culturation.

11. A serum suitable for treatment of swine infected with mystery swine disease, comprising:

the semi-purified blood serum of a mammal inoculated with an infectious agent derived from swine tissue infected with mystery swine disease.

12. A serum according to claim 11 wherein the infectious agent is obtained from an inoculum of a filtered homogenate of swine lung tissue infected with mystery swine disease.

13. A serum according to claim 11 wherein the infectious agent is obtained from the cell culture medium of in vitro cultured, avian, insect, or mammalian organ cells treated with an inoculum obtained from swine lung tissue infected with mystery swine disease.

14. A serum according to claim 13 wherein said cultured mammalian organ cells are a simian cell line.

15. A method for diagnosis of mystery swine disease in swine, comprising:

obtaining a lung tissue sample from the swine,  
forming a liquid homogenate of the sample,

-53-

adding the liquid homogenate to a vessel for immobilizing a viral material to form an immobilized mixture,

adding to the immobilized mixture a nonswine, mammalian species antibody serum to mystery swine disease to form a complex,

adding a labeled anti-species antibody to detect the complex.

16. A method according to claim 15 wherein the nonswine mammalian species antibody is a monoclonal antibody.

17. A method according to claim 15 wherein the labeled anti-species antibody carries a radioactive or color producing enzyme label.

18. A method according to claim 16 wherein said monoclonal antibody is SDOW 12 or SDOW 17.

19. A method for diagnosis of mystery swine disease in swine, comprising:

forming a liquid homogenate of a sample suspected of harboring the mystery swine disease infectious agent tissue sample from the swine,

adding to the liquid homogenate an immunoreactive antibody which is immunospecific for mystery swine disease virus,

detecting the presence of a precipitate of an antibody-antigen complex.

20. A method according to claim 19 wherein said antibody is SDOW 12.

21. A method according to claim 19 wherein said antibody is SDOW 17.

-54-

22. A method of making a MSD vaccine containing an inactivated or attenuated MSD infectious agent, the method comprising:

identifying swine lung tissue having thickened aveolar septae, degenerating cells and debris in aveolar spaces;

homogenizing the swine lung tissue with a pharmaceutically acceptable aqueous solution to form a mixture;

filtering the mixture through a series of filters to produce a filtered homogenate containing the MSD infectious agent; and

inactivating or attenuating the MSD infectious agent in the filtered homogenate to produce the MSD vaccine.

23. A method according to claim 22, wherein the homogenate is filtered to contain particles having a size no greater than about 1.0 micron.

24. A method according to claim 22 wherein the size is no greater than about 0.5 microns.

25. A method according to claim 22 wherein the size is no greater than about 0.1 microns.

26. A biologically pure culture of a fastidious, non-hemagglutinating, enveloped RNA virus, said culture being capable of affecting swine infertility and respiratory disease in swine, together with all zoopathogenic mutants thereof.

27. A monoclonal antibody produced by a murine derived hybrid cell line wherein said antibody is capable of specifically binding to at least one antigenic determinant of a fastidious non-hemagglutinating,

-55-

enveloped RNA virus, capable of affecting swine infertility and respiratory disease in swine.

28. A monoclonal antibody according to claim 27 wherein said antibody is an immunoglobulin molecule of the type IgG or IgM.

29. A monoclonal antibody according to claim 27 designated SDOW 12.

30. A monoclonal antibody according to claim 27 designated SDOW 17.

31. A biologically pure culture of swine infertility and respiratory syndrome virus, ATCC VR-2332, said culture being capable of affecting swine infertility and respiratory disease in swine, together with all zoopathogenic mutants thereof.

32. A vaccine composition comprising modified, live swine infertility and respiratory syndrome virus, ATCC VR-2332, and a pharmaceutically acceptable carrier, which virus has been modified to render the virus non-zoopathogenic in swine.

33. A vaccine composition comprising killed or inactivated swine infertility and respiratory syndrome, ATCC VR-2332, and a pharmaceutically acceptable carrier.

34. A method of immunizing swine against swine infertility and respiratory syndrome which comprises administering to swine a vaccine composition as recited in claim 32.


35. A method of immunizing swine against swine infertility and respiratory syndrome which comprises

administering to swine a vaccine composition as recited in claim 33.


36. A method of growing and isolating swine infertility and respiratory syndrome virus, ATCC VR-2332, which comprises inoculating the virus on a full or partial sheet of simian cells in the presence of serum in a suitable growth medium and incubating the inoculated cell sheet at about 34°C to 37°C until CPE is observed.

37. The method as recited in claim 36 wherein the simian cell line is MA-104.


**FIG. 1A**




**FIG. 1B**



**FIG. 2A**




**FIG. 2B**



3/5

FIG. 3



SUBSTITUTE SHEET

**FIG. 4A**



**FIG. 4B**





FIG. 5



# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 92/06873

<b>I. CLASSIFICATION OF SUBJECT MATTER</b> (If several classification symbols apply, indicate all) *		
According to International Patent Classification (IPC) or to both National Classification and IPC		
IPC <sup>5</sup> : A 61 K 39/00, A 61 K 39/12, G 01 N 33/569, G 01 N 33/577, C 12 N 7/00, C 12 N 7/02, C 12 N 7/04		
<b>II. FIELDS SEARCHED</b>		
Minimum Documentation Searched <sup>7</sup>		
Classification System	Classification Symbols	
IPC <sup>5</sup>	A 61 K, G 01 N 33/00, C 12 N	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched <sup>8</sup>		
<b>III. DOCUMENTS CONSIDERED TO BE RELEVANT <sup>9</sup></b>		
Category <sup>9</sup>	Citation of Document, <sup>11</sup> with indication, where appropriate, of the relevant passages <sup>12</sup>	Relevant to Claim No. <sup>13</sup>
NO DOCUMENTS HAVE BEEN DISCLOSED		
<p>* Special categories of cited documents: <sup>14</sup></p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"I" document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"Δ" document member of the same patent family</p>		
<b>IV. CERTIFICATION</b>		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
11 November 1992	25 NOV 1992	
International Searching Authority	Signature of Authorized Officer	
EUROPEAN PATENT OFFICE	SCHNASS e.h.	